INDEX

Africa, South, plant-food consumption in, 75–6.
Ammonium phosphate, as fertilizer, 154–62, 239–50.
— — in Germany and Denmark, 158–9.
— — in Great Britain, 158, 239–50.
— — in New Zealand, 160.
— — in N. America, 159.
— — in the Tropics, 160.
— — on acid soils, 248–9.
Ammonium sulphate, effect of, on sugar-cane, with lime, 83–8.
— — for coffee in S. India, 307.
— — for hill pastures in Wales, 58–63.
— — for sugar-cane in Queensland, 23.
— — production of, in Empire countries, 68.
Ammophos, 159–60.
Askew, H. O., *see* Rigg, T.
Australia, composition of natural pastures in, 236–7.
— consumption of N, P, K fertilizers in, 71.
— grassland improvement in, 228 sqq.
Australia, Queensland, Bureau of Sugar Experiment Stations, 20.
— sugar-cane in, 20–8.
— — manurial experiments on, 21–4.

B. radicicola, effect of, on wild white clover and grasses, 331–2.
Barley, effect of sulphuric-acid spray, on yield, 226; on malting quality, 221–2.
Basic slag, experiments with, on grasses and clovers, 324 sqq.
— — production of, in Canada, 69.
Birdsfoot trefoil in agriculture, 274–83.
— — cultivation for fodder, 275; for seed, 281.
Blackman, G. E., weed-control in cereal crops by sulphuric acid, 213–27.
Bones and bone-products in the Empire, 69.
Bush-sickness, soil and mineral supplements in treatment of, 1–8.
Buttercup, corn, control of, 218–19.

Cacao, branch systems, 49; disease, 49; root-systems, 48; seedling stocks, 48; transmission of yielding-power, 49–50.
— propagation by budding, 40–1, 47; cuttings, 43–5; grafting, 40, 47; layering, 45–7, 49.
Canada, basic slag production in, 69.
— experiments on wheat-quality in, 119–38.
— imports of fertilizers into, 72–3.
Canary-grass, use of, in nutritional experiments, 339–40.
Ceylon, imports of fertilizers into, 77.
Charlock, white, control of, 217–19.
— yellow, control of, 214–17.
Cheesman, E. E., vegetative propagation of cacao, 40–50.
Chippendale, H. G., *see* Davies, R. O.
Clover, wild white, effect of *B. radicicola* on development of, 331–2.
— effects of, on *Molinia* pasture, 330–1.
Clovers, effect of clipping on, 388.
— in Australia, 228 sqq.
— response of, to treatment on acidic upland soils, 324–36.
Coffea arabica, in S. India, 303 sqq.
— — root-system of, 293–302.
— — water-absorption by roots of, 301–2.
Coffee, diseases of, 192, 201, 303 sqq.
— pests of, 192, 200, 303 sqq.
— vegetative propagation of, 189–204; advantages and disadvantages of, 190–2; future of, 199; methods of, 189; practice of, 192–8.
Coffee-planting in S. India, 303–14.
— early work on, 305.
— work at Mysore Experimental Station, 305 sqq.; control of diseases and pests, 306, 307–11; manuring, 307; plant-breeding, 306–7, 311–13.
Coleman, L. C., coffee-planting in S. India, 303–14.
Coombs, A. V., border effect in plot experiments, 315–23.
Cotton, green-manuring experiments with, 101.
Craig, N., influence of maturity and rainfall on properties of lateritic soils in Mauritius, 349–58.
Cramer, P. J. S., early experiments on grafting coffee in Java, 200–4.
Crampton, E. W., nutritive value of pasture herbage, 337–48.

Dairy herd, changes and events in large, 139–52.
— births, sex-ratio, and heifers reared, 140–2.
— cows in, abortion, 149–51.
— — — and dead at birth, 143–4.
— — — and milk-yields, 151–2.
— — age of, in herd and drafted out, 144 sqq.
— — calving record, 137.
— — life of, 152.
— herd policy, 139.
— management, 139.
— wastage, 142–3, 147–9.
Davies, R. O., response of grasses, &c., to treatment on acidic upland soils, 324–36.

Davies, Wm., pasture legumes in relation to grassland improvement in Australia, 228–38.
Davies, W. Morley, soil survey of N. Shropshire, 178–88, 359–79.
Disease, bush-sickness, 1–8.
— canker of cacao, 49.
— *Hemileia* of coffee, 192, 201, 303, 306–8.
Diseases, of coffee, 192, 201, 303 sqq.
— of sugar-cane, 27–8.

Faulkner, O. T., experiments with leguminous crops in S. Nigeria, 93–102.
Fertilizers, and the Empire, 65–78.
— concentrated complete, experiments with, 239–50.
— effect of, on hill pastures, 58–63.
— — on yield of grasses, &c., 332–4.
Ferwerda, F. P., vegetative propagation of coffee, 189–99.
Fish manure for sugar-cane, 112, 117–18.
Fraser, A. H. H., wool-growth in sheep as affected by carbohydrate in diet, 9–19.

Grass, feeding value of fertilized and unfertilized, 345–7.
Grasses, effect of early defoliation on, 388–9.
— in hill pastures, relative palatability of, 51–8.
— on acidic upland soils, response to treatment of, 324–36.
Gray, A. N., Fertilizers and the Empire, 65–77.
Great Britain and Ireland, trade and consumption of fertilizers, 70.
Green-manure crops, burning of, in S. Nigeria, 97 sqq.
— for sugar-cane soils in Queensland, 23.
— residual value of, 94, 100.
Green-manuring, experiments in S. Nigeria, 93–102.
Guano, sources of, 67.

Halais, P., *see* Craig, N.
Heath, O. V. S., apparatus for measuring soil compactness in field, 205–12.
Hill, A. Glendon, sugar-cane breeding, 285–92.
Hudson, A. W., rate of seeding of wheat in relation to variety trials, 29–39.
Hunter-Smith, J., changes and events in a large dairy herd from 1912 to 1932, 139–53.

India, British, production of potassium nitrate in, 68.
Iron ammonium citrate as remedy for bush-sickness, 2 sqq.
Irrigation, effect of, on wheat soils and N-content of wheat in Canada, 123–7, 129–31.

Java, grafting of coffee in, 189–99.
— — — early experiments on, 200–4.

Kerr, H. W., sugar-cane in Queensland, 20–8.

Legumes, pasture, in Australia, 228–38.
Leguminous crops, experiments with, in S. Nigeria, 93–102.
Lewis, A. H., manurial effectiveness of ammonium phosphate, 154–62, 239–50.
Lime, interaction of, with ammonium sulphate in sugar-cane soil, 85–7.
— — — effect of potash and phosphate on, 89–90.
Limestone, use of, on acidic upland soils, 324 sqq.
Lime-requirement of sugar-cane soils, 79 sqq.
— measurement of, 91.
Liming, effect of, on sugar-cane yields, 78–92.
— in presence of ammonium sulphate, 83–7.
— on exhausted soils, 82–3.
— on peat-mosses, 169.
Limonite ore as remedy for bush-sickness, 2 sqq.
Lucerne, effect of clipping in first season, 381–3.
— effect of cutting on drought-resistance, 387–8.
— effect of cutting on thickness of stand, 385–7.
— effect of cutting on yield, &c., 380–90.
— effect of number of cuttings on later yields, 383–5.
— in Australia, 234 sqq.

Maize, yields of, after *Mucuna*, 95 sqq.
Maize-starch, effect on wool-growth in sheep, 10 sqq.
Mangolds, ammonium phosphate for, 239 sqq.
Marrow-stem kale, edge-row experiments with, 315 sqq.
Mauritius, lateritic soils in, 349–58.
Milton, W. E. J., relative palatability of seeds-mixtures, and influence of fertilizers on natural hill-pastures, 51–64.
Molasses, sugar-cane, manurial value of, 24–5.
Molinia, pastures, improvement of, by manuring and firing, 59–61.
— soil, response of herbage plants to treatment on, 324–36.
Moorland, reclamation of, 163–73.
— by *Fehncultur*, 166.
— by paring and burning, 166.
— in Scotland, 166–7, 172–3.
— modern methods of, 167 sqq.; cropping, 171; cultivation, 170–1; drainage, 167–9; liming, 169; manuring, 170.
Mucuna utilis, experiments with, in S. Nigeria, 94 sqq.

New Zealand, bush-sickness and allied ailments in, 1–8.

New Zealand, N, P, and K fertilizers, imports, &c., 74–5.
Nichols, J. E., *see* Fraser, A. H. H.
Nicol, Hugh, effect of cutting on yield, &c., of lucerne, 380–90.
Nitro-chalk, for acidic upland soils, 325 sqq.
Nitrophoska, use of, as fertilizer, 158–60, 240.
Nutman, F. J., root-system of *Coffea arabica*, 293–302.

Oats, ammonium phosphate for, 239 sqq.
— breeding of, in Scotland, 251–8; for early maturity, 253; heredity, 252; hybridization, 251; new varieties, 255–7; resistance to lodging, &c., 253–4.
— effect of sulphuric-acid spray on yield of, 226.
Ogg, W. G., reclamation of moorland, 163–73.
Owen, G., *see* Davies, W. M.

Palatability of seeds-mixtures, 51–64.
Pasture herbage, nutritive value of, 337–48.
Pastures, Australian, composition of, 236–7.
Pastures, hill, effect of fertilizers on, 58–63.
Pen manure, preparation and composition of, 103–4.
— — for sugar-cane, 104–9.
— — as affected by soil moisture, 112.
— — effects on yield, 113–15.
— — — with lime, 115–16.
— — optimum dressing of, 106–8.
— — value of N-content, 111–12.
— — *v.* inorganic manures, 109–12.
— — *v.* other organic manures, 117–18.
Pests, of coffee, 192, 200, 303 sqq.
— of sugar-cane, 20, 27–8, 291.
Phaseolus lunatus, experiments with, in S. Nigeria, 99–100.
Phosphate rock, Empire sources of, 66–7.
Pigs, nutritional anaemia in, 4–5.
Plant-food, consumption of, in Great Britain and Ireland, 70; in S. Africa, 75–6.
Poppy, field, control of, 219–20.
Potash, effect of, on yield of sugar-cane, 114.
— for acidic upland soils, 325 sqq.
Potassium nitrate, production of, in India, 68.
Potatoes, ammonium phosphate for, 239 sqq.
Protein in grassland herbage, quality of, 337 sqq.

Rigg, T., soil and mineral supplements in treatment of bush-sickness, 1–8.
Robb, Wm., oat-breeding in Scotland, 251–7.
Robertson, Ian M., *see* Ogg, W. G.
Robinson, D. H., birdsfoot trefoil in agriculture, 274–83.
Robinson, G. W., the soils of Wales, 258–73.

Sampling-observations, observer's bias in, 174–7.
Scotland, oat-breeding in, 251–7.
— reclamation of moorland in, 166–7, 172–3.
Seeding rate of wheat, 29–39.
— effect of, on quality, size, and yield of grain, 34 sqq.
Seeds-mixtures (grass), palatability of, 51–8.
— — effect of lime and fertilizers on, 59.
Sheep, bush-sickness in, 1–8.
— palatability of timothy and red clover to, 53 sqq.
— wool-growth in, as affected by carbohydrate in diet, 9–19.
Sheep-manure for sugar-cane, 117.
Shropshire, North, geography and climate, 178–9; geology, 179–84; soil survey of, 178–88, 359–79.
Shutt, F. T., quality of wheat as influenced by environment, 119–38.
Sinclair, K. J., *see* Lewis, A. H.
Soil compactness, apparatus for measuring, 205–7.
— conditions, effect of, on protein in wheat, 119–38.
— of Nelson, New Zealand, for bush-sickness, 2 sqq.
— surveys in Queensland, 26.
Soils, acid, ammonium phosphate on, 248–9.
— — upland, response of grasses, &c., on, 324–36.
— lateritic in Mauritius, 349–58.
— of N. Shropshire, 184–8, 359–79.
— of Wales, 262–73; brown earths, 266–9; dune, 270–1; impeded drainage, 271–2; limestone, 269–70; peat 272; podsols, 263–6; skeletal, 266.
— sugar-cane, in Queensland, 25–6; in Trinidad, 78–92, 103–18.
— wheat, in Canada, 'breaking' and cultivated, 121–2.
— — irrigated and non-irrigated, 123–6.
Stafford, W. C., *see* Hudson, A. W.
Sugar-beet, ammonium phosphate for, 239 sqq.
— edge-row experiments with, 315 sqq.
Sugar-cane, breeding of, 285–92; field-trial problems, 289–90; for resistance to pests, 290–1; history of, 285–6, 291; technique of, 287–9.
— quality of, due to liming, 92.
— yield of, with lime, 78–92; with pen manure, 104–9; with phosphate and potash, 89–90.
— in Queensland, 20–8.
— — pests of, 20.
— — plant-breeding and pest-control, 27–8.
— — soils, 25–6.
— — testing maturity of, 26–7.
— — utilization of by-products, 24–5.
Sulphur, Empire sources of, 65.

Sulphuric acid, use of, for weed control, 213–27.
— effect on germinative capacity of charlock, 215.
— — quality of malting barley, 221–2.
Superphosphate, Empire production of, 69.
— for coffee in S. India, 307.
— for grassland in Australia, 236.
— for hill pastures in Wales, 58–63.
— for sugar-cane in Queensland, 23.

Timothy grass, use of, in nutritional experiments, 339–40.
Trefoil, birdsfoot, 274–83.
Trefoils, in Australia, 231.
Trevains, D., *see* Lewis, A. H.
Trinidad, experiments on sugar-cane in, 78–92, 103–18.
Turner, P. E., sugar-cane and sugar-cane soils, 78–92, 103–18.

Variety trials, seeding-rate of wheat in, 29–39.
Vegetative propagation, of cacao, 40–50.
— of coffee, 189–204.

Wales, climate of, 260; geology of, 259; soils of, 258–73.
Watson, D. J., *see* Yates, F.
Weeds in cereals, control of, with sulphuric acid, 213–27; charlock, 214–16; corn buttercup, 218–19; field poppy, 219–20; white charlock, 217–18.
— — effect on crop-yields, 216–20, 226.
Wheat, effect of sulphuric-acid spray on yield of, 226.
— effect of temperature, rainfall, and irrigation on quality of, 118–38.
— 'lodging' of, as affected by rate of seeding, 37.
— nitrogen-content of, in Canada, 122, 124–7, 129, 131.
— rate of seeding of, in variety trials, 29–39.
— sampling-observations on, 174–7.
— yields, as affected by grain drills, 38, 39.
Wheatland, Canada, 'breaking' and cultivated soils, 122 sqq.
— influence of irrigation, 123–7, 129–30.
Wool, sheep's, effect of dietetic carbohydrate on fleece-weight, 11; fibre-length, 13; fineness, 15; fibre-volume, 16; follicle-number, 17.

Yams, green-manuring experiments with, 101.
Yates, F., observer's bias in sampling-observations on wheat, 174–7.

ERRATUM

Vol. II, April 1934, p. 94, 10th line from base, *in lieu of* 0·5 *read* 5·0.